

Cleaning Products, Environmental Awareness and Risk Perception in Mérida, Mexico

Ruth Magnolia Martínez-Peña^{1*}, Almira L. Hoogesteijn², Stephen J. Rothenberg³, María Dolores Cervera-Montejano², Julia G. Pacheco-Ávila⁴

1 Independent researcher, 113 Woodview Terrace, Hurst, Texas, United States of America, **2** Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, CINVESTAV, Departamento de Ecología Humana, Unidad Mérida. Km. 6 Antigua Carretera a Progreso, Mérida, Yucatán, México, **3** Instituto Nacional de Salud Pública, INSP, Universidad No. 655 Colonia Santa María Ahucatlilán, Cerrada Los Pinos y Caminera, Cuernavaca, Morelos, México, **4** Universidad Autónoma de Yucatán, UADY, Facultad de Ingeniería, Av. Industrias no contaminantes por Periférico Norte Apartado Postal 150, Cordemex, Mérida, Yucatán, México"

Abstract

Cleaning products are associated with many health and environmental problems. Contamination of water resources by cleaning products is more likely to occur with septic tanks as sewage treatment systems especially in karstic terrains. We explored women's ideas about water sources and the risk cleaning products pose to health and sewage in Mérida, a city in the Yucatán peninsula of Mexico. Women were unaware of the city's water management system. We found a positive and statistically significant association between risk perception and environmental awareness, education level and employment status. We suggest developing education and risk communication strategies to inform residents about the hydro-geological features in the Yucatán, the vulnerability of its karstic aquifer and the health and environmental risks associated with cleaning agents.

Citation: Martínez-Peña RM, Hoogesteijn AL, Rothenberg SJ, Cervera-Montejano MD, Pacheco-Ávila JG (2013) Cleaning Products, Environmental Awareness and Risk Perception in Mérida, Mexico. PLoS ONE 8(8): e74352. doi:10.1371/journal.pone.0074352

Received May 16, 2013; **Accepted** August 5, 2013; **Published** August 23, 2013

Copyright: © 2013 Martínez-Peña et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Funding provided by Consejo Nacional de Ciencia y Tecnología (México). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: ruthmmp@hotmail.com

Introduction

Although cleaning has been a common activity in an historical context, it is since the 19th century that complex mass marketed cleaning products form part of our cleaning habits [1]. During the last century thousands of different cleaning products have been produced, generating phenomenal global growth in the industry [2]. Advertising of cleaning products has been mostly directed at women and is supported by the idea that cleaning is synonymous with happiness [3] and health. Nonetheless, studies have identified the adverse consequences of cleaning products on human health [4] and the environment [5]. Contact dermatitis [6], asthma [7] and other respiratory disorders [8], poisoning [9] and bacterial resistance [10] have been associated with the use of cleaning products. Endocrine disruption has also been associated with exposure to cleaning products [11]. Adverse consequences in the environment include eutrophication [12], anatomical and physiological changes in aquatic and terrestrial fauna [13], the elicitation of estrogenic response in mammals and fish [14], the facilitation of the adsorption of pharmaceutical compounds such as acetaminophen and carbamazepine by aquifer materials and sediments [15], and the creation of water-

repellent soils due to irrigation with gray water which contains surfactants [16].

Nearly 12% of the global land surface is formed by carbonate rocks, prone to chemical dissolution, i.e. karstification [17]. Karstic aquifers are the source of drinking water for about 25% of the world population [18], in both, developed [19] and developing countries [20]. Karstic aquifers are vulnerable to contamination [21], particularly from septic tank leachate [22]. In the United States, where about 20% of total households use septic tanks [23], boron [24] and phosphorus [25] derived from detergents were found in aquifers contaminated by septic waste in karstic terrain.

Popular awareness of the vulnerability of fresh water resources and concerns about the role played by the detergents in water pollution and the emergence of health problems can be explored through the concepts of Environmental Awareness (EA) and Risk Perception (RP). EA is the information that people have about any phenomena related to their environment [26,27], their concern about the environment [28], and their willingness to act in favor of the environment, including the behavior derived from that commitment [29]. The concept of EA has been used to analyze popular knowledge, concerns and attitudes about issues such

as household electronic devices [30], water pollution [31], conservation [32] and climate change [33]. The RP concept functions in the context of the evaluation of judgments about activities and technologies considered to be risky [34]. RP studies have analyzed popular knowledge, attitudes and behaviors related to water reuse [35], pesticides [36], loss of agro-biodiversity [37] and West Nile virus [38]. When the object of the RP study is a product, RP refers to how much that people feel that their safety is threatened by using it [39]. To our knowledge there are no published studies relating cleaning products, RP and EA in the developing world in general and Mexico in particular, and cleaning products contamination studies are scarce. Detergent compounds have been found in the San Juan River in the state of Nuevo León [40], in the Texcoco River in central Mexico [41] and in the Lerma-Santiago River in the state of Mexico [42]. In the Yucatán Peninsula, high concentrations of phosphates associated with household and hotel sewage have been found in the coastal lagoons of Chelem in Yucatán and in Nichupte and Bojórquez in Quintana Roo [43].

Mérida's karstic surface [44] was formed during the Pliocene-Miocene epoch [45]. The permeability of the karstic formation facilitates infiltration [46], increases groundwater vulnerability to pollution [47] and explains the lack surface water [48]. The aquifer is the only source of fresh water in the zone and its availability in the entire state is affected by saltwater intrusion, overexploitation [49] and by contamination [50]. In the urban area, water from the top 20 meters of the aquifer is not suitable for human consumption anymore [51]. As Mérida lacks a conventional sewerage system [52] untreated effluents infiltrate the groundwater [53]. Sewage management in Mérida relies upon septic tanks, most of which are not built to proper technical specifications [54,55] and thus hinder wastewater transformation, chemical degradation and biodegradation processes [56]. The aim of this paper is to analyze the answers of 739 women in Mérida, the capital city of the state of Yucatán (Mexico), which explored their knowledge of water sources and sewage treatment in the city and the effect of environmental awareness on perceived environmental and health risks related to household cleaning products. This study supplies information to risk communicators and health and environmental authorities applicable not only in Mérida, but in every city with karstic aquifers and on-site sewage disposal systems around the world.

Materials and Methods

Study site

Mérida is situated in the southeast of Mexico, between 20°41' N and 21°12' N, and 89°27' W and 89°49' W, at an altitude ranging from 7 and 10 m. a. s. l. [57]. Mérida has 830,732 inhabitants and covers approximately 88,300 ha (218,194 acres) [58].

Sampling design

Mothers of children attending any of the six levels of primary school were invited to participate. Two criteria led to the focus on this population: the emphasis of cleaning product

Table 1. Sociodemographic characteristics (control variables) of the study group.

Variable	Data included in OLS		Data excluded from OLS		p [*]
	N	%	N	%	
Age					0.920
≤ 29	133	19.6	6	16.2	
30–44	491	72.4	28	75.7	
≥ 45	54	8.0	3	8.1	
Education level ^{**}					0.604
Low	263	38.8	11	47.8	
Middle	142	20.9	5	21.7	
High	273	40.3	7	30.4	
Income ^{***}					0.965
≤ 4	221	39.6	10	43.5	
>4 – 10	111	19.9	5	21.7	
>10 – 25	177	31.7	7	30.4	
>25	49	8.8	1	4.4	
Occupation					0.629
Housewife	382	56.3	20	60.6	
Outside employee	296	43.7	13	39.4	

^{*} p-values were calculated from Fisher exact and chi-squared tests, depending on the observed value in the cells.

^{**} Low: no educated-secondary school; middle: incomplete high school-incomplete college; and high: complete college-postgraduate.

^{***} Income is expressed as multiples of the minimum monthly wage in 2007 (MXN \$ 1,428 or USD\$116.4). Income data were reported only by 581 participants.

advertisements on the role of women in cleaning activities [59], and the vulnerability of children under 12 to poisoning by cleaning products [60]. Six public schools and 14 private schools were selected based on geographic distribution (north, center and south of the city). This selection was important given the socioeconomic segregation of the city, in which people of high income live to the north site and people of low income live in the southern site [61]. A total of 739 surveys were completed, 294 in public schools and 445 in private schools. 599 surveys were answered in school facilities and 140 were sent to the respondents' homes and collected at the school one week later. We divided our survey in three sections: Environmental Awareness Scale (EAS, Table 1), Risk Perception Scale (RPS, Table 2) and sociodemographic data (age, education, income and occupation). EAS had 10 items with three possible answers: True, False and Not sure. RPS was a Likert scale with 16 items and five possible answers: Totally agree, Agree, Not sure, Disagree, and Totally disagree. A pilot version of the survey was conducted with 40 respondents, based on the results some changes were made to the final version. The surveys were performed between August and November 2007.

The research project and the techniques to obtain the data were approved by the Professors Board of the Human Ecology Department of CINVESTAV, according to Postgraduate Studies Regulations (Art. 43). In each school, a verbal authorization to send each child home with a written invitation for the mother to participate was provided by the general director. That invitation had the information about the date and

Table 2. Rasch Model Fitness diagnoses for the Environmental Awareness Scale (EAS).

ITEM*	Difficulty (logits)	Standard error	Outfit		Answers (%)	
			MNSQZ		Right	Wrong
A septic tank is a container that transforms gray water into clean water.	-1.84	0.12	0.82	-1.2	86	2
The sludge from septic tanks is obtained and transformed into fertilizer.	0.88	0.09	0.83	-2.9	42	10
The ocean is the source of tap water.	-0.57	0.09	0.87	-1.9	68	7
The water extracted from wells comes from underground water sources.	-0.56	0.09	0.95	-0.7	69	9
Almost the 100% of gray water in the city leaks into the ground and can contaminate the underground sources of water.	-0.94	0.1	0.97	-0.3	74	6
The gray water from households is transformed into drinking water in treatment plants.	0.78	0.09	1.01	0.2	43	18
Rain is the source of tap water.	-0.67	0.09	1.04	0.5	70	5
The gray water from septic tanks goes into the ground.	-0.01	0.09	1.13	2.1	59	13
Tap water comes from underground water sources.	0.45	0.09	1.08	1.4	51	17
Sludge and gray water from septic tanks go to a hole on the ground.	2.48	0.11	1.33	2.1	17	30

* Items are listed in order of MNSQ.

the place where the survey would be conducted. It also stated the voluntary nature of this activity and the confidentiality to manage personal data. Once in the room, the women who attended the invitation were asked again about their intention to participate, and pictures of this process were taken. The fact that the women responded to the invitation and gave verbal recognition of their intent to participate confirmed their consent. Following the request of the directors of three of the private schools, the survey was attached to the invitation to give the women the opportunity to complete it at home. In this case, the consent was indicated by returning the completed survey.

Statistical Analysis

The Rasch model [62] was used in order to process the ordinal data from EAS and RPS as continuous data, all assumptions were met. Rasch analysis is a probabilistic model based on the assumption that the measuring scale measures only one dimension or latent variable (EA or RP in our case) which can be used to rate the item’s difficulty as well as the subject’s ability [63]. Subjects and items share the same measurement scale in logits or the log-odds transformation of the probability of a response [64]. The scale to calibrate the difficulty of the items has zero (0) as the middle point, which means an equal likelihood to answer it correctly or incorrectly [65]. Zero is also the middle point of the scale to calibrate the ability of the subjects and means an equal likelihood to make a mistake or to have a correct answer [66]. The probability of a right or expected answer is a function of the difference between the difficulty of the item and the ability of the subject [67]. Unlike the classic approaches to measure abilities which emphasize fitting models to data, Rasch analysis emphasizes fitting data to the model [68]. Under the Rasch model, an item provides information about a subject’s performance when the quadratic mean of Outfit, MNSQ (an outlier sensitive fit statistic that picks up rare events), is between 0.5 and 1.5 [69].

Fitting of EAS and RPS to the Rasch model was analyzed with *Winsteps* software [70]. Fifteen surveys were omitted due to a lack of answers to the EAS items. All the EAS items (10) had MNSQ values between 0.5 and 1.5. The EA score was obtained for 724 participants. In the RPS analysis the original five possible answers were reduced to three by grouping Totally Agree and Agree into the Agree category, and Totally Disagree and Disagree into the Disagree category. Twenty-three surveys were not considered due to a total lack of answers in the RPS items. Two items had Outfit MNSQ higher than 1.5 and were removed from the final analysis of difficulty and ability. The RP score was obtained for 716 participants.

Prior to analysis, data were tested for normality using Skewness/Kurtosis, Shapiro–Wilk and Shapiro–Francia tests. Student’s t-tests were performed to compare EA and RP scores between groups with and without complete data in the sociodemographic section of the survey. To analyze the relationship between the dependent variable EA score and independent variables OLS regression analysis was performed, only with surveys containing complete data on age, education and occupation (n = 678). Independent variables considered were age (≤ 29, 30–44 and ≥ 45), education level (low: no educated–secondary school; middle: incomplete high school–incomplete college; and high: complete college–postgraduate) and occupation (housewife and outside employee). A standard errors adjustment was performed to avoid heteroscedasticity effects. To analyze the relationship between the dependent variable RP score and independent variables (EA score, age, education level and occupation) OLS regression analysis was performed. Income was considered a potential control variable in both models (EA and RP) but it was not included due to collinearity and low response rate. OLS regression was performed with a standard errors adjustment and 10000 bootstrap repetitions to avoid heteroscedasticity effects and the lack of residual normality.

The data were processed with *Access 2002* and *Excel 2002* [71], *Stata 10 2007* [72] and *Winsteps* [73].

Results

No statistically significant differences were found between surveys completed at the school versus those completed at home. No significant differences were found between sociodemographic characteristics of participants who were included and those who were excluded from the OLS regression analysis (Table 1). The maximum and minimum difficulty values for the EAS items were 2.48 and -1.84, respectively, with zero as mean and standard deviation (SD) of 1.14 (Table 2). The maximum and minimum EA ability values were 4.16 and -3.97, respectively, with zero as mean and SD of 1.28. The mean of EA scores by sociodemographic variable and level is shown in Figure 1. The percentages of right and wrong answers to each EAS item are shown in Table 2.

About 70% of the participants were aware that rain and the ocean are not sources of tap water; 51% knew that the actual water source is groundwater (Table 2). More than 70% of the participants were not aware about the environmental fate of sludge and gray water; 43% of participants knew that the local treatment systems do not turn sewage water into drinkable water, and 42% knew that sludge is not used by the local authorities to produce fertilizers (Table 2). More than 70% of the participants were aware that the aquifer is vulnerable to contamination (Table 2).

Two items of the RPS were removed from the analysis due to their high difficulty values (5.61 y 3.65, respectively). The maximum and minimum difficulty values for the 14 items retained in the RPS were 1.35 and -3.22, respectively, with zero as mean and SD of 1.19 (Table 3). The maximum and minimum RP ability values were 4.27 and -1.58, respectively, with zero as mean and SD of 1.32. The mean of RP scores by sociodemographic variable and level is shown in Figure 2. The practices of reading the product labels as well as keeping the products out of the reach of children were related to a higher RP (Table 3). The practices of mixing some cleaning products as well as using a greater amount of them were related to a lower RP (Table 3).

The OLS regression analysis for EA as dependent variable had an R² of 0.0849 (F_{5,678} = 14.38, p < 0.001). Age and education had a positive impact on the EA score (Table 4). The OLS regression analysis for RP as dependent variable had an R² of 0.2125 (Wald = 204.65, p < 0.01). Higher EA and education, as well as outside employment had positive impacts on RP score (Table 5). There was a significant difference between middle and high education odds ratio (Wald= 4.32, p < 0.05). Women who attended high school and college are twice as likely to have a higher RP as women without formal education or with only intermediate education (Table 5). A greater difference was found between participants with low education and those who attended college (odds ratio = 2.59) (Table 5). The RP scores by sociodemographic variables and levels are shown in the Figure 2.

Discussion

Table 3. Rasch Model Fitness diagnoses for the Risk Perception Scale (RPS).

ITEM*	Difficulty (logits)	Standard error	Outfit	
			MNSQZ	
Waste of cleaning products helps marine fauna.	-0.29	0.12	0.74	-2.2
Waste of cleaning products can contaminate groundwater.	-0.11	0.12	0.80	-1.8
Cleaning products are harmless.	0.11	0.11	0.88	-1.2
Waste of detergents helps to clean groundwater.	0.84	0.10	0.90	-1.6
It is possible to get clean surfaces using less detergent.	0.47	0.10	0.95	-0.6
It is necessary to use more detergent to get cleaner surfaces.	1.33	0.09	0.96	-0.8
It is important to read the cleaning products' labels.	-3.22	0.39	0.99	0.1
Mixing different cleaning products is good to clean better.	1.35	0.09	1.01	0.2
Cleaning products can affect people's health.	-0.07	0.12	1.07	0.7
Mixing cleaning products can be dangerous.	0.13	0.11	1.08	0.9
It is useless to read the cleaning product directions for use.	0.9	0.10	1.10	1.6
Waste of cleaning products reaches the ocean and affects marine fauna.	0.93	0.10	1.14	2.2
Cleaning products must be kept out of the reach of children.	-1.69	0.20	1.23	0.9
Children can ingest cleaning products by accident and get poisoned.	-0.66	0.14	1.39	2.3
Antibacterial detergents are better because they prevent diseases.	REMOVED			
Antibacterial detergents help some bacteria to get stronger.	REMOVED			

* Items are listed in order of MNSQ. Last two items were removed due to their MNSQ values were higher than 1.5.

Unlike previous studies on EA and RP that used Principal Component Analysis [74,75] and Factor Analysis [76–78], this study used Rasch Analysis [79]. Rasch Analysis allows for transforming ordinal data into continuous data, hence performing more robust statistical processes like regression analysis. However, our experience in this study confirmed what has been pointed out by previous researches [80] about the need to test Rasch Analysis during the survey design phase in order to guarantee its fitness to the model prior to application.

The EAS dealt with three issues: the sources of tap water, sewage management and vulnerability of groundwater to contamination. Only 51% of the participants knew that the actual source of tap water is groundwater (Table 2). What accounts for the remaining 49%? There are two possible answers. From a methodological perspective, it could be that half of the participants did not understand the concept of groundwater, which implies a bad item design. From a




Figure 1. Mean of Environmental Awareness (EA) in logits, according to the sociodemographic variables. A. Age; B. Education level (low: no educated–secondary school; middle: incomplete high school–incomplete college; and high: complete college–postgraduate); C. Income (expressed as multiples of Minimum Monthly Wage, MMW. One MMW = USD \$110); and D. Occupation.

doi: 10.1371/journal.pone.0074352.g001

functional perspective, it is possible that the conceptual link between turning on the tap and extracting water from a well is hard to establish. It would mean that the groundwater concept is easier linked to a well than to a tap.

Regarding sewage management, even when women have a general knowledge about septic tanks, they mostly ignore the process that occurs there. This result could be related to low

frequency of septic tank cleaning—even when water authorities suggest an annual cleaning [81]—or with a lack of information provided by the cleaning companies. Only 18% of participants thought that treatment systems turn sewage water into drinkable water, and 10% thought that sludge is used to produce fertilizers (Table 2). These low percentages are encouraging since they indicate that most of women are aware




Figure 2. Mean of Risk Perception (RP) in logits, according to the sociodemographic variables. A. Age; **B.** Education level (low: no educated–secondary school; middle: incomplete high school–incomplete college; and high: complete college –postgraduate); **C.** Income (expressed as multiples of Minimum Monthly Wage, MMW. One MMW = USD \$116); and **D.** Occupation.

doi: 10.1371/journal.pone.0074352.g002

of the substandard sewage management in place, it also indicates a need for clear and true information about it.

The high percentage of women knowing that the aquifer is vulnerable to contamination could be explained by the increasing public concern about the environment [82] rather than from knowledge of the hydro-geo-ecological features of a karst system. This concern was expressed globally in the Brundtland Report of 1987 and with the United Nations

initiative Decade of Education for Sustainable Development, 2005-2014. Around the time the data were collected, there were initiatives that included aquifer contamination as an issue of public concern [83,84]. The scope of this research does not analyze the impact of these local programs.

Almost two decades ago, Ronald Inglehart pointed out that the increasing concern for environmental issues was related to a change in the priorities of the societies, moving from a

Table 4. Results of OLS regression analysis for Environmental Awareness (EA) Scores as dependent variable (n = 678; R² = 0.0849).

Environmental Awareness [*]	Coefficient ^{**}	t	p	Confidence interval (95%) ^{***}
Age (30–44) ^a	1.54	3.54	< 0.001	1.21–1.95
Age (≥45) ^a	1.93	3.33	0.001	1.31–2.86
Middle Education ^b	1.21	1.36	0.173	0.91–1.61
High Education ^b	1.66	3.74	< 0.001	1.27–2.16
Outside employee ^c	1.12	0.99	0.321	0.89–1.40
Constant	-0.79	-2.31	0.021	0.64–0.96

^{*} Environmental Awareness was measured in logits or the log-odds transformation of the probability of a response.

^{**} Regression coefficients are presented as exp(logits).

^{***} Confidence intervals were calculated using heteroscedasticity-correct standard errors.

^a Comparison group: < 29 years.

^b Comparison group: Low Education

^c Comparison group: Housewife.

Table 5. Results of OLS regression analysis for Risk Perception (RP) Scores as dependent variable (n = 678; R² = 0.2125).

Risk Perception [*]	Coefficient ^{**}	z	p	Confidence interval (95%) ^{***}
Environmental Awareness Score	1.24	4.40	0.000	1.13–1.36
Age (30–44) ^a	1.23	1.68	0.092	0.97–1.57
Age (≥45) ^a	1.35	0.15	0.881	0.69–1.53
Middle Education ^b	2.00	5.48	0.000	1.56–2.58
High Education ^b	2.59	7.46	0.000	2.02–3.33
Employee	1.24	2.07	0.038	1.01–1.53
Constant	3.63	12.33	0.000	2.96–4.46

^{*} Risk Perception was measured in logits or the log-odds transformation of the probability of a response.

^{**} Regression coefficients are presented as exp(logits).

^{***} Confidence intervals were calculated using heteroscedasticity-correct standard errors.

^a Comparison group: < 29 years.

^b Comparison group: Low Education.

^c Comparison group: Housewife.

materialistic point of view to a post materialistic perspective, where meeting basic needs implies acquiring new concerns [85]. Detractors of Inglehart’s ideas maintain that higher environmental awareness is not related to higher wealth but to higher vulnerability to environmental hazards [86,87]. In this

study, women with higher education obtained higher EA and RP scores (Table 4). Regardless of Inglehart’s assumption, it is likely that our results are related to better access to information and more diverse sources of social contact than with a different matrix of values. This situation could also explain the higher EA and RP scores in outside employees and women older than 29. It is important to take into account that more years of education mean more opportunities to develop the skills useful in responding surveys such as the one analyzed here.

Even when the global results of EAS show the overall effect of older age and more education, a closer examination of the results shows a lack of information about specific issues i.e. the disposal of sludge and gray water. This situation should be considered by environmental authorities and encourage the design of environmental education activities, which need to take into account the cultural diversity. The city could take School Education Boards to spread information to the public via the Schools.

Regarding the RPS, the diagnoses of its fitness to the Rasch Model forced us to remove two items, both related to increases in bacterial resistance associated with antibacterial cleaning products (Table 3), due to their high difficulty values. From the methodological perspective, it is necessary to rephrase those items to be useful in future studies. Notwithstanding, it is important to consider the impact of advertising that presents antibacterial products as a means of disease prevention and well-being [88]. Once those two items were removed, the items related to mixing cleaning products and the need to use more of each product to get better results, had the highest difficulty values (Table 3). Results show that the mixing of cleaning products, e.g. chlorine and hydrochloric acid is not perceived as a risk [89], and as our results show not related to a high RP score. By contrast, reading product labels and keeping the products out of the reach of children are practices related to a higher RP. Nonetheless, the fact of having a higher RP does not ensure that people actually perform these practices [90]. Studies have shown that most consumers do not read the cleaning product labels [91] or they think that those directions are exaggerated taking into account the product quality [92]. In our particular case, most of the cleaning products are bottled or bagged in small containers without labels; this is a common practice in small retailer stores. Store owners buy in bulk and retail small quantities to maximize gain [93]. Whether the labels are incomplete, useless, ignored or entirely missing, the environmental and health outcomes are the same: an increase of cleaning products in the sewage system, a considerable number of women affected by chlorine fumes and many children exposed to dangerous products.

Some studies about environmental risk perception or pro-environmental behavior supports the idea that young people with higher education and better health have a higher pro-environmental attitude [94], more perception about contamination, and are more willing to participate in pro-environmental activities [95]. Other studies point out that the related factor of lower earning opportunity does not necessarily mean a lower environmental RP [96] because the vulnerability of poor communities to environmental risks makes them more aware of this context [97]. Regarding health risks, higher

education does not equate to a higher RP [98]. Despite the opposite conclusions of the cited studies, it is a fact that sociodemographic variables have an impact on RP [99], even when the results vary depending on the research purposes and techniques [100].

Our results show that more education and a formal employment have a positive impact on RP (Table 5). This result is consistent with previous studies [101]. However, as we said regarding the EAS results, it is necessary to consider the relationship between years of education and improved skills to answer questionnaires.

The last analysis we performed was to examine the link between EA and RP. According to our results, knowledge about water sources and sewage management in the city has a positive relationship with the cleaning product risk perception. This means that enhancing EA might improve RP and thus modify use patterns involving cleaning products. This would imply that less cleaning agents would pass to the aquifer and women and children's exposure would be diminished. While the results indicate that a key to modifying behavior could be the implementation of educational strategies that increase knowledge about water sources and sewage treatment systems in the city of Mérida, a risk we propose three main ideas as communication strategy: a) the dangers of mixing cleaning products; b) the benefits of using fewer types of cleaning products; c) the benefits of using less product and achieve the same results. These ideas address the health and environmental risks derived from exaggerated cleaning products use. Even when the messages are simple, the principal obstacle to such a campaign would be that it opposes commercial interests [102] and deeply rooted practices linked to the use of products women are very familiar with [103].

References

- Quitau M (2004) Changing ideas of bodily cleanliness. International Summer Academy on Technology Studies. Urban Infrastructure in Transition. Available: <http://orprints.org/3298/1/3298.pdf>. Accessed 2007 December 21.
- ACNielsen (2006) Mercados en crecimiento alrededor del mundo: productos para el hogar. Available: <http://www.acnielsen.com.mx/press/documents/MercadosenCrecimiento-ProductosparaelHogar.pdf>. Accessed 2007 January 10.
- Shrikhande V (2003) Stereotyping of women in television advertisements. Masters dissertation, University of Louisiana. Available: http://etd.lsu.edu/docs/available/etd-0516103-141609/unrestricted/Shrikhande_thesis.pdf. Accessed 2007 September 14.
- Adelson L, Sunshine I (1952) Fatal poisoning due to a cationic detergent of the quaternary ammonium compound type. *Am J Clin Pathol* 22: 656-661. PubMed: 14933349.
- Zahn RK, Zahn G, Müller WE, Müller I, Beyer R et al. (1977) Consequences of detergent pollution of the sea: effects on regenerating sponge cubes of *Geodia cydonium*. *Sci Total Environ* 8: 109-151. doi:10.1016/0048-9697(77)90072-9. PubMed: 905819.
- Lim YL, Goon A (2007) Occupational skin diseases in Singapore 2003-2004: an epidemiologic update. *Contact Dermatitis* 56: 157-159. doi:10.1111/j.1600-0536.2007.01032.x. PubMed: 17295692.
- Zock JP, Vizcaya D, Le Moual N (2010) Update on asthma and cleaners. *Curr Opin Allergy Clin Immunol* 10: 114-120. PubMed: 20093933.
- Bello A, Quinn MM, Perry MJ, Milton DK (2010) Quantitative assessment of airborne exposures generated during common cleaning tasks: a pilot study. *Environ Health* 9: 76. doi:10.1186/1476-069X-9-76. PubMed: 21118559.
- Williams H, Moyns E, Bateman DN, Thomas SH, Thompson JP et al. (2012) Hazard of household cleaning products: A study undertaken by the UK National Poisons Information Service. *Clin Toxicol (Phila)* 50: 770-775. doi:10.3109/15563650.2012.709936. PubMed: 22809171.
- Aiello AE, Marshall B, Levy SB, Della-Latta P, Lin SX et al. (2005) Antibacterial cleaning products and drug resistance. *Emerg Infect Dis* 11: 1565-1570. doi:10.3201/eid1110.041276. PubMed: 16318697.
- Zota AR, Aschengrau A, Rudel RA, Brody JG (2010) Self-reported chemicals exposure, beliefs about disease causation, and risk of breast cancer in the Cape Cod Breast Cancer and Environment Study: a case-control study. *Environ Health* 9: 40. doi:10.1186/1476-069X-9-40. PubMed: 20646273.
- Al-Rawi SM (2005) Contribution of man-made activities to the pollution of the Tigris within Mosul area/IRAQ. *Int J Env Res Pub He* 2: 245-250. PubMed: 16705824.
- Falconer IR (2006) Are endocrine disrupting compounds a health risk in drinking water? *Int J Env Res Pub He* 3: 180-184. PubMed: 16823090.
- Zoller U (2006) Estuarine and coastal zone marine pollution by the nonionic alkylphenol ethoxylates endocrine disrupters: is there a potential ecotoxicological problem? *Environ Int* 32: 269-272. doi:10.1016/j.envint.2005.08.023. PubMed: 16225920.
- Hari AC, Paruchuri RA, Sabatini DA, Kibbey TC (2005) Effects of pH and cationic and nonionic surfactants on the adsorption of pharmaceuticals to a natural aquifer material. *Environ Sci Technol* 39: 2592-2598. doi:10.1021/es048992m. PubMed: 15884354.
- Wiel-Shafran A, Ronen Z, Weisbrod N, Adar E, Gross A (2006) Potential changes in soil properties following irrigation with surfactant-rich greywater. *Ecol Eng* 26: 348-354. doi:10.1016/j.ecoleng.2005.12.008.
- Ford DC, Williams PW (2007) Karst hydrogeology and geomorphology. Chichester: Wiley. 562pp.
- Ford DC, Williams PW (2007) Karst hydrogeology and geomorphology. Chichester: Wiley. 562pp.
- Goldscheider N (2012) A holistic approach to groundwater protection and ecosystem services in karst terrains. *Aqua Mundi* 3: 117-124.

Taking these factors into account, the effectiveness of risk communication activities could depend on the participation of all the stakeholders in the community [104] as well as on the testimonies of affected people [105].

Our results show the necessity to implement environmental education and risk communication. Hydro-geo-ecological conditions in Mérida will not change and septic tanks will continue to be used. This is also true for millions of households found in karst areas in the United States [106], France [107], Australia [108] and many other regions around the world [109]. The risk of contamination to groundwater by cleaning products will not decrease unless people change daily practices. Even though acquiring new information does not immediately lead to new behaviors, it is important to begin educating the community about water resources and sewage management. Increasing community awareness about the vulnerability of their water resources as well as the health risks associated with cleaning products is the first step to develop healthier and environmentally friendly practices and spaces.

Acknowledgements

RMMP thanks Mike Linacre for his guidance in utilizing *Winsteps*, and Michael R. Miller and David Buitrago Tello for their editing assistance.

Author Contributions

Conceived and designed the experiments: ALH RMMP. Performed the experiments: RMMP. Analyzed the data: SJR RMMP ALH MDCM JGPA. Contributed reagents/materials/analysis tools: SJR. Wrote the manuscript: RMMP.

20. León LM, Parise M (2008) Managing environmental problems in Cuban karstic aquifers. *Environ Geol* 58: 275-283.
21. Bakalowicz M (2005) Karst groundwater: a challenge for new resources. *Hydrogeol J* 13: 148-160. doi:10.1007/s10040-004-0402-9.
22. Pacheco AJ, Cabrera SA (1997) Groundwater contamination by nitrates in the Yucatán Peninsula, Mexico. *Hydrogeol J* 5: 47-53. doi: 10.1007/s100400050113.
23. EPA (2009) Septic Systems Fact Sheet. Available: http://water.epa.gov/aboutow/owm/upload/2009_06_22_septics_septic_systems_factsheet.pdf. Accessed 2013 June 25.
24. Verstraeten I, Fetterman G, Meyer M, Bullen T, Sebree S (2005) Use of tracers and isotopes to evaluate vulnerability of water in domestic wells to septic waste. *Ground Water Monit Rev* 25: 107-117. doi:10.1111/j.1745-6592.2005.0015.x.
25. McCray JE, Kirkland SL, Siegrist RL, Thyne GD (2005) Model parameters for simulating fate and transport of on-site wastewater nutrients. *Ground Water* 43: 628-639. doi:10.1111/j.1745-6584.2005.00777.x. PubMed: 16029188.
26. Langford IH (2002) An existential approach to risk perception. *Risk Anal* 22: 101-120. doi:10.1111/0272-4332.t01-1-00009. PubMed: 12017353.
27. Lima ML, Barnett J, Vala J (2005) Risk perception and technological development at a societal level. *Risk Anal* 25: 1229-1239. doi: 10.1111/j.1539-6924.2005.00664.x. PubMed: 16297227.
28. Varkuti A, Kovacs K, Stenger-Kovacs C, Padisak J (2008) Environmental awareness of the permanent inhabitants of towns and villages on the shores of Lake Balaton with special reference to issues related to global climate change. *Hydrobiologia* 599: 249-257. doi: 10.1007/s10750-007-9190-2.
29. Zsoka A (2008) Consistency and "awareness gaps" in the environmental behavior of Hungarian companies. *J Clean Prod* 16: 322-329.
30. Huang P, Zhang X, Deng X (2006) Survey and analysis of public environmental awareness and performance in Ningbo, China: a case study on household electrical and electronic equipment. *J Clean Prod* 14: 1635-1643. doi:10.1016/j.jclepro.2006.02.006.
31. Anderson B, Romani A, Phillips H, Wentzel M, Tabela K (2007) Exploring environmental perceptions, behaviors and awareness: water and water pollution in South Africa. *Popul Environ* 28: 133-161.
32. Ballantyne R, Packer J, Hughes K (2008) Environmental awareness, interests and motives of botanic gardens visitors: implications for interpretive practice. *Tourism Manage* 29: 439-444. doi:10.1016/j.tourman.2007.05.006.
33. Varkuti A, Kovacs K, Stenger-Kovacs C, Padisak J (2008) Environmental awareness of the permanent inhabitants of towns and villages on the shores of Lake Balaton with special reference to issues related to global climate change. *Hydrobiologia* 599: 249-257. doi: 10.1007/s10750-007-9190-2.
34. Slovic P (1987) Perception of Risk. *Science* 236: 200-285. doi:10.1126/science.3494308. PubMed: 3494308.
35. Baggett S, Jeffrey P, Jefferson B (2006) Risk perception in participatory planning for water reuse. *Desalination* 187: 149-158. doi:10.1016/j.desal.2005.04.075.
36. Crane M, Norton A, Leaman J, Chalak A, Bailey A et al. (2006) Acceptability of pesticide impacts on the environment: what do United Kingdom stakeholders and the public value? *Pest Manag Sci* 62: 5-19. doi:10.1002/ps.1134. PubMed: 16261541.
37. Schmidt MR, Wei W (2006) Loss of agro-biodiversity, uncertainty & perceived control: a comparative risk perception study in Austria and China. *Risk Anal* 26: 455-470. doi:10.1111/j.1539-6924.2006.00744.x. PubMed: 16573633.
38. Zielinski-Gutierrez E, Hayden M (2006) A model for defining West Nile virus risk perception based on ecology and proximity. *Ecohealth* 3: 28-34. doi:10.1007/s10393-005-0001-9.
39. Ortiz J, Resnick M, Kengskool K (2000) The effects of familiarity and risk perception on workplace warning and compliance. In: XIVth Triennial Congress of the International Ergonomics Association and 44th Annual Meeting of the Human Santa: Factors and Ergonomics Society Mónica, California
40. Flores-Laureano J, Nívar J (2002) An assessment of stream water quality of the Rio San Juan, Nuevo León, Mexico, 1995-1996. *J Environ Qual* 31: 1256-1265. doi:10.2134/jeq2002.1256. PubMed: 12175044.
41. Belmont MA, Ikonoumou M, Metcalfe CD (2006) Presence of nonylphenol ethoxylate surfactants in a watershed in central Mexico and removal from domestic sewage in a treatment wetland. *Environ Toxicol Chem* 25: 29-35. doi:10.1897/04-648R.1. PubMed: 16494221.
42. Pineda-Flores G, Hernández T, Cruz MC, Gutiérrez-Castrejón T (1999) Application of 2 microbioassays for evaluating the pollution present in the Xochimilco and Lerma-Santiago basins. *Revi Latinoam Microbiol* 41: 251-258. PubMed: 11001710.
43. Herrera-Silveira J (2006) Lagunas costeras de Yucatán (SE, México): investigación, diagnóstico y manejo. *Ecotropicos* 19: 94-108.
44. Steinich B, Velázquez-Oliman G, Marín L, Perry E (1996) Determination of the ground water divide in the karst aquifer of Yucatán, Mexico, combining geochemical and hydrogeological. *Geofis Int* 35: 153-159.
45. Perry E, Swift J, Gamboa J, Reeve A, Sanborn R et al. (1989) Geologic and environmental aspects of surface cementation, north coast, Yucatán, Mexico. *Geology* 17: 818-821. doi: 10.1130/0091-7613(1989)017.
46. Batllori-Sampedro E, González-Piedra J, Díaz-Sosa J, Febles-Patrón J (2006) Caracterización hidrológica de la región Costera noroccidental del estado de Yucatán, México. *Investigaciones Geográficas* 59: 74-92.
47. Pacheco J, Cabrera A, Pérez R (2004) Diagnóstico de la calidad del agua subterránea en los sistemas municipales de abastecimiento en el estado de Yucatán, México. *Ing Rev Académica* 8: 165-179.
48. Escolero O, Marín L, Steinich B, Pacheco A, Cabrera S et al. (2002) Development of a protection strategy of karst limestone aquifers: the Mérida, Yucatán, Mexico case study. *Water Resour Manag* 16: 351-367. doi:10.1023/A:1021967909293.
49. Escolero O, Marín L, Steinich B, Pacheco J, Molina-Maldonado A et al. (2005) Geochemistry of the hydrogeological reserve of Mérida, Yucatán, Mexico. *Geofis Int* 44: 301-314.
50. Graniel E, Vera I, González L (2004) Dinámica de la interfase salina y calidad del agua en la Costa Nororiental de Yucatán. *Ing Rev Académica* 8: 15-25.
51. Escolero O, Marín L, Steinich B, Pacheco J, Molina-Maldonado A et al. (2005) Geochemistry of the hydrogeological reserve of Mérida, Yucatán, Mexico. *Geofis Int* 44: 301-314.
52. Quintal C (2003) Problemática de la disposición de aguas residuales en la ciudad de Mérida. In: Tello. p. L; Alonzo A, editor *Evolución y estrategias del desarrollo urbano ambiental en la Península de Yucatán*. pp. 207-215
53. Batllori-Sampedro E, Febles-Patrón J (2002) El agua subterránea en el desarrollo regional de la península de Yucatán. *Av Perspectiva* 21: 67-77.
54. Magaña A (2002) Reactor anaerobio horizontal doable (RAH-D) su desarrollo en la FIUADY. *Ing Rev Académica* 6: 61-65.
55. Febles-Patrón J, Hoogsteijn A (2008) Análisis del marco legal para la protección del agua subterránea en Mérida, Yucatán. *Ing Rev Académica* 12: 71-79.
56. McCray JE, Kirkland SL, Siegrist RL, Thyne GD (2005) Model parameters for simulating fate and transport of on-site wastewater nutrients. *Ground Water* 43: 628-639. doi:10.1111/j.1745-6584.2005.00777.x. PubMed: 16029188.
57. Nacional de Estadística y Geografía Instituto INEGI (2009) Prontuario de información geográfica municipal de los Estados Unidos Mexicanos. Available: <http://mapserver.inegi.org.mx/dsis/prontuario/index2.cfm>. Accessed 2013 January 30
58. Nacional de Estadística y Geografía Instituto INEGI (2010) INEGI. Censo de Población y Vivienda. Cuestionario básico. Available: <http://www3.inegi.org.mx/sistemas/mexicocifras/default.aspx?src=487&e=31>. Accessed 2013 January 30
59. Habib RR, El-Masri A, Heath RL (2006) Women's strategies for handling household detergents. *Environ Res* 101: 184-194. PubMed: 16616916.
60. Forrester MB (2006) Patterns of exposures at school among children age 6-19 years reported to Texas Poison Centers, 1998-2002. *J Toxicol Env Heal A* 69: 263-268. PubMed: 16407086.
61. Cruz M (2009) Diagnóstico sobre la realidad social, económica y cultural de los entornos locales para el diseño de intervenciones en materia de prevención y erradicación de la violencia en la Región sur: el caso de la zona metropolitana de Mérida, Yucatán. Comisión Nacional para Prevenir y Erradicar la Violencia contra las Mujeres. Secretaría de Gobernación. México.
62. Wright B, Stone M (1999) *Measurement essentials*. Wilmington. Wide Range Inc.. p. 221.
63. Wright B, Stone M (1999) *Measurement essentials*. Wilmington. Wide Range Inc.. p. 221.
64. Wright B, Stone M (1999) *Measurement essentials*. Wilmington. Wide Range Inc.. p. 221.
65. Wright B, Stone M (1999) *Measurement essentials*. Wilmington. Wide Range Inc.. p. 221.
66. Wright B, Stone M (1999) *Measurement essentials*. Wilmington. Wide Range Inc.. p. 221.
67. Wright B, Stone M (1999) *Measurement essentials*. Wilmington. Wide Range Inc.. p. 221.

68. Engelhard G (1984) Thorndike, Thurstone and Rasch: a comparison of their methods of scaling psychological and educational tests. *Appl Psychol Meas* 8: 21-38.
69. Linacre M (2012) A User's Guide to Winsteps. Available: <http://www.winsteps.com/a/winsteps-manual.pdf>. Accessed 2012 September 1.
70. Linacre M (2007) Winsteps. USA: Chicago.
71. Microsoft (2002). Corporation. Redmond, USA: Office.
72. StataCorp LP (2007) Stata 10. USA: College Station.
73. Linacre M (2007) Winsteps. USA: Chicago.
74. Schmidt MR, Wei W (2006) Loss of agro-biodiversity, uncertainty and perceived control: a comparative risk perception study in Austria and China. *Risk Anal* 26: 455-470. PubMed: 16573633.
75. Willis HH, Dekay ML (2007) The roles of group membership, beliefs and norms in ecological risk perception. *Risk Anal* 27: 1365-1380. PubMed: 18076502.
76. Lemyre L, Lee J, Mercier P, Bouchard L, Krewski D (2006) The structure of Canadians' health risk perceptions: environmental, therapeutic and social health risks. *Health Risk Soc* 8: 185-195.
77. Slimak MW, Dietz T (2006) Personal values, beliefs and ecological risk perception. *Risk Anal* 26: 1689-1705. PubMed: 17184406.
78. Toma L, Mathijs E (2007) Environmental risk perception, environmental concern and propensity to participate in organic farming programs. *J Environ Manage* 83: 145-157. PubMed: 16697103.
79. Wright B, Stone M (1999) Measurement essentials. Wilmington. Wide Range Inc. p. 221.
80. Tennant A, McKenna SP, Hagell P (2004) Application of Rasch Analysis in the development and application of quality of life instruments. *Value Health* 7: S22-S26. PubMed: 15367240.
81. Comisión Nacional del Agua CNA (1997) Norma Oficial Mexicana NOM-006-CNA-1997, Fosas sépticas prefabricadas, especificaciones y métodos de prueba. Available: <http://www.profepa.gob.mx/innovaportal/file/3302/1/nom-006-conagua-1997.pdf>. Accessed 2007 October 17.
82. Franzen A (2003) Environmental attitudes in international comparison: an analysis of the ISSP surveys 1993 and 2000. *Soc Sci Q* 84: 297-308.
83. de Mérida Ayuntamiento (2007). Plan Municipal Desarrollo: 2007-2010. Available: <http://201.159.134.38/listPoder4.php?edo=31&orderSeleccionado=2324&catTipo=2324>. Accessed 2007 October 17
84. de Desarrollo Secretaría Urbano y Medio Ambiente SEDUMA (2008); Ambiental Cultura. Available: http://www.seduma.yucatan.gob.mx/main.php?s=s_cultura. Accessed 2008 April 19
85. Anderson B, Romani A, Phillips H, Wentzel M, Tabela K (2007) Exploring environmental perceptions, behaviors and awareness: water and water pollution in South Africa. *Popul Environ* 28: 133-161.
86. Drori I, Yuchtman-Yaar E (2002) Environmental vulnerability in public perceptions and attitudes: the case of Israel's urban centers. *Soc Sci Q* 83: 53-63.
87. White M, Hunter L (2005) Public perceptions of environmental issues in a developing setting. University of Colorado Research Program on Environment and Behavior. Working Paper EB2005-003 Available: <http://www.colorado.edu/ibs/pubs/eb/eb2005-0003.pdf>. Accessed 2007 December 13.
88. Aiello A, Larson E, Levy S (2007) Consumer antibacterial soaps: effective or just risky? *Clin Infect Dis* 45: 137-147.
89. Gorguner M, Aslan S, Inandi T, Cakir Z (2004) Reactive airways dysfunction syndrome in housewives due to a bleach-hydrochloric acid mixture. *Inhal Toxicol* 16: 87-91. PubMed: 15204781.
90. Zsoka A (2008) Consistency and "awareness gaps" in the environmental behavior of Hungarian companies. *J Clean Prod* 16: 322-329.
91. Kovacs DC, Small MJ, Davidson CI, Fischhoff B (1997) Behavioral factors affecting exposure potential for household cleaning products. *J Expo Anal Environ Epidemiol* 7: 505-520. PubMed: 9306234.
92. Habib RR, El-Masri A, Heath RL (2006) Women's strategies for handling household detergents. *Environ Res* 101: 184-194. PubMed: 16616916.
93. Martínez-Peña RM (2008) Percepción de riesgo del uso de productos de limpieza domésticos en Mérida, Yucatán. Thesis to obtain a Master's Degree in Sciences. Centro de Investigación y Estudios Avanzados del Instituto Politécnico Nacional, CINVESTAV-Unidad Mérida. Departamento de Ecología Humana..
94. Wakefield SE, Elliott SJ, Eyles JD, Cole DC (2006) Taking environmental action: the role of local composition, context and collective. *Environ Manage* 37: 40-53. PubMed: 16328677.
95. Toma L, Mathijs E (2007) Environmental risk perception, environmental concern and propensity to participate in organic farming programs. *J Environ Manage* 83: 145-157. PubMed: 16697103.
96. Tilt B (2006) Perceptions of risk from industrial pollution in China: a comparison of occupational groups. *Hum Organ* 65: 115-127.
97. Slimak MW, Dietz T (2006) Personal values, beliefs and ecological risk perception. *Risk Anal* 26: 1689-1705. PubMed: 17184406.
98. Williams A, Lindsell C, Rue L, Blomkalns A (2007) Emergency Department education improves patient knowledge of coronary artery disease risk factors but not the accuracy of their own risk perception. *Prev Med* 44: 520-525. PubMed: 17336373.
99. Wakefield SE, Elliott SJ, Eyles JD, Cole DC (2006) Taking environmental action: the role of local composition, context and collective. *Environ Manage* 37: 40-53. PubMed: 16328677.
100. Visschers V, Meertens R, Passchier W, De Vries N (2007) An associative approach to risk perception: measuring the effects of risk communications directly and indirectly. *J Risk Res* 10: 371-383.
101. Wakefield SE, Elliott SJ, Eyles JD, Cole DC (2006) Taking environmental action: The role of local composition, context and collective. *Environ Manage* 37: 40-53. PubMed: 16328677.
102. ACNielsen (2006) Mercados en crecimiento alrededor del mundo: productos para el hogar. Available: <http://www.acnielsen.com.mx/press/documents/MercadosenCrecimiento-ProductosparaelHogar.pdf>. Accessed 2007 January 10.
103. Ortiz J, Resnick M, Kengskool K (2000) The effects of familiarity and risk perception on workplace warning and compliance. Proceedings of the International Ergonomics Association. Annual Meeting of Santa: The Human Factors and Ergonomics Society Mónica, California
104. Peres F, Moreira JC, Rodrigues KM, Claudio L (2006) Risk perception and communication regarding pesticide use in rural work: a case study in Rio de Janeiro State, Brazil. *Int J Occup Env Heal* 12: 400-407. PubMed: 17168229.
105. de Wit JB, Das E, Vet R (2008) What works best: objective statistics or a personal testimonial? An assessment of the persuasive effects of different types of message evidence on risk perception. *Health Psychol* 27: 110-115. PubMed: 18230021.
106. Katz BG, Griffin DW, McMahon PB, Harden HS, Wade E et al. (2010) Fate of effluent-borne contaminants beneath septic tank drainfields overlying a karst aquifer. *J Environ Qual* 39: 1181-1195. PubMed: 20830905.
107. Vaute L, Drogue C, Garrelly L, Ghelfenstein M (1997) Relations between the structure of storage and the transport of chemical compounds in karstic aquifers. *J Hydrol* 199: 221-238.
108. Levett KJ, Vanderzalm JL, Page DW, Dillon PJ (2010) Factors affecting the performance and risks to human health of on-site wastewater treatment systems. *Water Sci Technol* 62: 1499-1509. PubMed: 20935366.
109. Goldscheider N (2012) A holistic approach to groundwater protection and ecosystem services in karst terrains. *Aqua Mundi* 3: 117-124.